








Complications With Demineralized Bone Matrix, Hydroxyapatite and Beta-Tricalcium Phosphate in Single and Two-Level Anterior Cervical Discectomy and Fusion Surgery

Global Spine Journal
2024, Vol. 14(2S) 78S–85S
© The Author(s) 2023
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: [10.1177/21925682231157320](https://doi.org/10.1177/21925682231157320)
journals.sagepub.com/home/gsj


Juan P. Cabrera, MD^{1,2} , Sathish Muthu, MS^{3,4} , Mohamed Kamal Mesregah, MD⁵ , Ricardo Rodrigues-Pinto, MD, PhD⁶ , Neha Agarwal, PhD⁷, Viswanadha Arun-Kumar, MBBS, MS⁸, Yabin Wu, PhD⁹ , Gianluca Vadalà, MD, PhD¹⁰, Christopher Martin, MD¹¹, Jeffrey C. Wang, MD¹², Hans Jörg Meisel, MD, PhD⁷ , Zorica Buser, PhD^{13,14} , and AO Spine Knowledge Forum Degenerative

Abstract

Study Design: Systematic literature review.

Objectives: To analyze the evidence available reporting complications in single or two-level anterior cervical discectomy and fusion (ACDF) using a demineralized bone matrix (DBM), hydroxyapatite (HA), or beta-tricalcium phosphate (β -TCP).

Methods: A systematic review of the literature using PubMed, EMBASE, Cochrane Library, and [ClinicalTrials.gov](https://clinicaltrials.gov) databases was performed in August 2020 to identify studies reporting complications in one or two-level ACDF surgery using DBM, HA, or β -TCP. The study was reported following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.

Results: A total of 1857 patients were included, 981 male and 876 female, across 17 articles; 5 prospective, and 12 retrospective. We noted heterogeneity among the included studies concerning the study design and combination of graft materials utilized in them. However, we noted a higher incidence of adjacent segment disease (17.7%) and pseudoarthrosis (9.3%) in fusion constructs using DBM. Studies using β -TCP reported a higher incidence of pseudoarthrosis (28.2%) and implant failures (17.9%).

¹ Department of Neurosurgery, Hospital Clínico Regional de Concepción, Concepción, Chile

² Faculty of Medicine, University of Concepción, Concepción, Chile

³ Department of Orthopaedics, Government Medical College, Dindigul, India

⁴ Orthopaedic Research Group, Coimbatore, India

⁵ Department of Orthopaedic Surgery, Menoufia University Faculty of Medicine, Shebin El-Kom, Egypt

⁶ Spinal Unit/Unidade Vertebro-Medular (UVM), Department of Orthopaedics, Centro Hospitalar Universitário do Porto, Porto, Portugal

⁷ Department of Neurosurgery, BG Klinikum Bergmannstrost Halle, Germany

⁸ Reva Spine Centre, Visakhapatnam, AP, India

⁹ Research Department, AO Spine, AO Foundation, Davos, Switzerland

¹⁰ Campus Bio-Medico University of Rome and Fondazione Policlinico Universitario Campus Bio-Medico, Rome, Italy

¹¹ Department of Orthopaedic Surgery, University of Minnesota, Minneapolis, MN, USA

¹² USC Spine Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

¹³ Gerling Institute, Brooklyn, NY, USA

¹⁴ Department of Orthopaedic Surgery, NYU Grossman School of Medicine, New York, NY, USA

Corresponding Author:

Juan P. Cabrera, Department of Neurosurgery, Hospital Clínico Regional de Concepción, San Martín 1436, Concepción, Chile.

Email: jucabrera@udec.cl



Creative Commons Non Commercial No Derivs CC BY-NC-ND: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Conclusions: Degenerative cervical conditions treated with one or two-level ACDF surgery using DBM, HA, or β -TCP with or without cervical plating are associated with complications such as adjacent segment disease, dysphagia, and pseudarthrosis. However, consequent to the study designs and clinical heterogeneity of the studies, it is not possible to correlate these complications accurately with any specific graft material employed. Further well-designed prospective studies are needed to correctly know the related morbidity of each graft used for achieving fusion in ACDF.

Keywords

demineralized bone matrix, hydroxyapatite, beta-tricalcium phosphate, anterior cervical discectomy and fusion, complications, pseudarthrosis, dysphagia

Introduction

Single or two-level anterior cervical discectomy and fusion (ACDF) surgery is a surgical procedure with a low risk of complications.¹ However, there is significant variability among surgeons regarding the perception of impact and severity of reported complications in the literature. In fact, despite dysphagia being one of the most frequently observed adverse events in ACDF surgery, 58% of surgeons consider it as a minor complication with significant disagreement.² In the context of cervical degenerative diseases, the main aim is to achieve decompression of the neural structures with interbody fusion, and an autograft is the first and most widely used for this purpose.

Nonetheless, donor site complications such as pain and infection due to the harvesting procedure can have negative consequences,³ and therefore different osteobiologics have been used to maintain the fusion rates in the place of autografts. In addition, there is great interest in the spine surgery community to understand more about osteobiologics because of the higher cost involved in their use. The commercially available grafts apart from the bone morphogenetic proteins (BMP) have emerged rapidly while their efficacy and complications with their use are not being completely understood.

In a constant effort to avoid complications in ACDF surgery, the commonly employed graft alternatives to autograft include substances such as demineralized bone matrix (DBM), hydroxyapatite (HA), and beta-tricalcium phosphate (β -TCP) to achieve promising results in the fusion rates during one or two-level ACDF surgery. The specific spectrum of related complications with their usage is lacking, which precludes the decision-making process in clinical practice for their routine usage. Therefore, this systematic review aims to analyze the best evidence available reporting complications in 1-2 level ACDF using these osteobiologics.

Methods

We performed a systematic review of the literature using PubMed, EMBASE, Cochrane Library, and ClinicalTrials.gov databases in August 2020 to identify studies reporting complications in one or two-level ACDF surgery using DBM, HA,

and/or β -TCP. The systematic review was reported following the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.⁴

Search Strategy, Inclusion, and Exclusion Criteria

Three authors (J.P.C., N.A, Y.W.) performed the database search and initially screened articles published from January 2000 to August 2020. The screening process of all articles was performed at the titles and abstracts level, and the selected articles were analyzed for eligibility in full-text and assessed by 2 independent reviewers. A mismatch in the selection of any article was resolved by a third reviewer. The inclusion and exclusion criteria are summarized in the PICO table ([Supplementary Material Table 1](#)).

Complications Included

We categorized the reported complications as follows: peri-operative surgical complications, medical complications, mid or long-term outcomes, and other complications.

Peri-Operative Surgical Complications. Neurologic deterioration ≥ 1 motor grade in American Spinal Cord Injury Association (ASIA)⁵ motor scale, new post-operative radiculitis or neuropathic pain, wound dehiscence, superficial wound infection, deep wound infection, airway compromise requiring surgical intervention, dysphagia, dysphonia/hoarseness of voice, anterior cervical hematoma/seroma, epidural hematoma/seroma, cerebrospinal fluid leak/meningocele, vertebral artery injury, esophageal perforation, Homer's Syndrome, and peri-operative death.

Mid or Long-Term Complications. Nonunion, construct failure with loss of correction (screw pull out, plate dislodgement), construct failure without loss of correction (screw pull out, plate dislodgement), and revision surgery for incomplete neurologic decompression and persistent symptoms.

Medical Complications. Cardiac arrest/failure/arrhythmia, deep vein thrombosis, delirium, gastrointestinal bleeding, myocardial

infarction, pneumonia, pressure sores, pulmonary embolism, systemic infection, and urinary tract infection.

Other Complications. Complications that were not included above.

List of Osteobiologics Included

Each study included in this systematic review was categorized according to the number of treated levels and the osteobiologics reported in the article. The osteobiologics included were demineralized bone matrix (DBM), hydroxyapatite (HA); beta-tricalcium phosphate (β -TCP).

Risk of Bias Assessment

We used the Cochrane tool (RoB-2)⁶ for the assessment of the risk of bias for randomized trials. In addition, non-randomized studies were assessed with the Methodological Index for Nonrandomized Studies (MINORS),⁷ for comparative and non-comparative studies. The maximum score is 16 for non-comparative studies and 24 for comparative studies.

Results

A total of 756 articles were found in the initial search of all four databases used. After the removal of duplicated

articles, 584 articles were analyzed and 431 of them were excluded. A total of 153 articles were assessed by two independent reviewers for eligibility and finally 17 articles – 5 prospective and 12 retrospective – were included.^{8–24} The search mechanism used in the review is shown in Figure 1.

There were 1857 patients included in the study with 981 male and 876 female patients. Single-level surgery was performed in 1231 patients, two-level ACDF surgery in 481 patients, and 145 patients were not differentiated between one or two levels. The minimum mean time of follow-up was 12 months. The demographics and other baseline characteristics are summarized in Table 1.

Risk of Bias Assessment

The perceived risk of bias across the retrospective studies was assessed using the MINORS assessment tool in twelve studies which included 5 non-comparative studies^{9,11–13,24} that scored from 8 to 12 points for a total of 16 points for non-comparative studies, and 7 comparative studies^{10,14,15,17,18,21,23} that scored from 15 to 21 points for a total of 24 points for comparative studies (Supplementary Material Table 2). The risk of bias across the prospective studies was assessed using RoB-2 in 5 randomized controlled trials, having 4 studies^{8,19,20,22} with a low risk of bias and 1 study¹⁶ with a high risk of bias (Figure 2).




Figure 1. PRISMA flow diagram for identification and selection of studies for analysis.

Overall Reported Complications

The most frequently reported complications among 1857 patients included from 17 articles were radiological and/or clinical adjacent segment disease (ASD) in 53 (2.85%) patients, followed by dysphagia/dysphonia or swallowing problems in 49 (2.64%) patients. We noted heterogeneity among the included studies concerning the study design and combination of graft materials utilized in them as shown in [Supplementary Material Table 3](#). However, we noted a higher incidence of adjacent segment disease (17.7%) and pseudoarthrosis (9.3%) in fusion constructs using DBM. Studies using β -TCP reported a higher incidence of pseudoarthrosis (28.2%) and implant failures (17.9%). We noted comparable dysphagia/dysphonia rates across all the osteobiologics analyzed. Individual risks of these complications among

osteobiologics such as DBM, HA, and β -TCP are provided as a heatmap in [Table 2](#).

Discussion

The present systematic review attempted to analyze the rate of postoperative complications associated with the use of DBM, HA, or β -TCP in ACDF surgery which resulted in identifying only 17 studies with suboptimal quality. From the included studies the most frequently reported complications to include adjacent segment disease, dysphagia, and pseudoarthrosis. Although we attempted to individualize the complications per the graft materials analyzed the heterogeneity of the included studies precluded us from generating strong conclusions for clinical practice related to safety.

Table 1. Demographic and Baseline Characteristics of the Included Studies.

First Author	Year	Country	Study Design	Number of Patients	Mean Age (Years)	Male/Female	1 Level ACDF	2 Levels ACDF
Hacker RJ et al ⁶	2000	USA	Prospective	488	44 ± 9/47 ± 7 [∞] 43 ± 8/47 ± 7 [∞] 44 ± 9/48 ± 8 [∞]	95/84 80/87 74/68	390	98
Agrillo U et al ⁷	2002	Italy	Retrospective	45	49.7	26/19	33	12
Cauthen JC et al ⁸	2003	USA	Retrospective	88	49.5 (34.4-66.1) 51.1 (32.9-67.4) 51.4 (29.6-70.3)	15/15 12/20 13/13	43	45
Cosar M et al ⁹	2008	Turkey	Retrospective	17	46 (33-60)	8/9	17	—
Akula M et al ¹⁰	2008	UK	Retrospective	50	51 (31-90)	31/19	42	8
Moon HJ et al ¹¹	2010	South Korea	Retrospective	27	50.8 (31-67)	18/9	—	27
Niu CC et al ¹²	2010	Taiwan	Retrospective	53	49.5 ± 11.5 52.2 ± 10.5	15/13 12/13	35	18
Zagra A et al ¹³	2013	Italy	Retrospective	86	32-60 28-29 29-63	13/11 15/14 17/16	80	6
Kim CH et al ¹⁴	2013	South Korea	Prospective	52	49.5 ± 12.8 55.2 ± 12.4	17/12 14/9	52	—
Shin JS et al ¹⁵	2014	Korea	Retrospective	60	50.0 ± 12.0 49.0 ± 11.0 44.3 ± 9.7	7/13 12/8 13/7	60	—
Kim SJ et al ¹⁶	2014	Korea	Retrospective	96	51.9 ± 13.5 46.5 ± 10.5	25/23 24/24	96	—
Xie Y et al ¹⁷	2015	China	Prospective	68	54.6 ± 12.7 56.3 ± 10.5	20/15 16/17	29	39
Yi J et al ¹⁸	2015	Korea	Prospective	77	51.9 ± 11.7 51.3 ± 12.4	25/13 26/13	77	—
Chin K et al ¹⁹	2017	United States	Retrospective	145	53 ± 1.0 53.4 ± 1.6	33/42 34/36	NR	NR
Farrokhi MR et al ²⁰	2017	Iran	Prospective	64	47.6 45.3	12/20 14/18	64	—
Yu J et al ²¹	2017	Korea	Retrospective	247	53.7 ± 10.2 53.8 ± 10.9 51.7 ± 9.9	43/33 46/36 65/24	115	132
Shiban E et al ²²	2018	Germany	Retrospective	194	54	91/103	98	96

[∞] denotes 1-level/2-level.

ACDF – Anterior Cervical Discectomy and Fusion.




Figure 2. Risk of Bias analysis for the prospective randomized trials included in the review using RoB-2 assessment tool.

The gold standard graft used in ACDF surgery has been the autologous iliac bone graft demonstrating good results in achieving intervertebral body fusion. However, it is not free from complications. The harvesting procedure of the autograft leads to donor-site complications³ such as pain, paresthesia, infection, and hematoma formation. Apart from the donor site-related complications, autograft requires additional plate instrumentation with screws to avoid graft collapse and better restoration of the cervical lordosis, increasing the cost and the rate of dysphagia.²⁵ Therefore, we performed this systematic review for understanding the spectrum of complications related to some osteobiologics growingly used in ACDF surgery such as DBM, HA, and β -TCP.

Notably, we found seventeen studies reporting complications in 1-2 level ACDF using the osteobiologics of interest such as DBM, HA, or β -TCP. However, the heterogeneity in the study designs and heterogeneity in the graft material usage prevented us from generating strong conclusions concerning their complications. We did not include complications such as donor-site complications from the studies using autografts and subsidence from the studies using cages in our analysis as

mentioned in the exclusion criteria to avoid further heterogeneity in the generated results.

Interbody fusion is the final objective in ACDF, and the fusion rate could be variable due to the radiological method used, time of follow-up, and subjective criteria employed.^{26,27} There was one study reporting fusion rates of 76% at 6 months of follow-up when a stand-alone PEEK cage filled with hydroxyapatite compared with 92% in tricortical iliac bone graft with plating.¹⁶ Nonetheless, hydroxyapatite showed over 87% of fusion rates when follow-up is until 12 months and mixed with autologous graft⁸ or with DBM/ β -TCP.²⁰ The negative face of the same outcome, ie non-union or pseudarthrosis, was reported in some of the included studies with heterogeneity. For example, Yi et al²⁰ compared a plate augmented PEEK cage construct filled with HA + DBM vs HA + β -TCP and reported that the rate of non-union at 12 months was 13% in both groups. Such combined usage of osteobiologic prevented us from identifying the material that is responsible for pseudoarthrosis. Moreover, the study of Yu et al²³ comparing stand-alone PEEK cage filled with DBM vs PEEK cage + DBM with plating vs iliac bone graft with plating reported

Table 2. Heatmap of the Complications Analysed Across the Included Studies.

Complications	Autograft	DBM	β -TCP	HA
Neck pain	0.6	2	NA	4.7
Dysphagia	4.8	4.2	NA	3
Dysphonia	2.9	3.1	4.7	2.4
ASD	2.8	17.7	3.1	NA
Pseudoarthrosis	2.5	9.3	28.2	1.2
Implant failure	0.9	1.8	17.9	0
Graft migration	3	0	0	0
Medical complications	1.4	3.8	0	0.6
Neurological worsening	0	2	0	2.3
Additional surgery	5.3	0	2.2	2.4
Infection	0	3.3	0	3.4
Dural tear	0.6	NA	NA	0

ASD: adjacent segment disease; DBM: demineralised bone matrix; β -TCP: beta tricalcium phosphate; HA: hyaluronic acid; NA: not available.

similar rates of fusion. Still, the supplementation with plate fixation especially while using autologous graft is beneficial in reducing the time to fusion. A recent study including 1-3 levels of ACDF treated with self-locking stand-alone cages filled with hydroxyapatite reported pseudoarthrosis at single level ACDF in 15.4% of patients while at 2-level ACDF in 15.2% of patients, and they reported that symptomatic pseudoarthrosis occurred in 6.1% requiring reoperation.²⁸

Dysphagia is one of the most common observed complications in ACDF and has significant heterogeneity in its reporting due to the variability in its definition and measuring methods.²⁹ Similarly, we noted a variable incidence of dysphagia across the included studies. Patients using autograft in their fusion construct reported the highest incidence of dysphagia (4.8%) followed by patients with DBM (4.2%) in their fusion construct. This could be explained by the higher use of plated constructs in the autograft group which resulted in a higher incidence of dysphagia.²⁵ Shin et al¹⁷ confirmed this phenomenon by comparing zero-profile and stand-alone PEEK cages packed with DBM vs iliac bone graft with plating with a significantly higher rate of dysphagia in the last group. However, cervical plating plays a role in preventing cage subsidence and restoration of cervical lordosis when compared with stand-alone cages.²⁵ Moreover, the incidence rates were comparable across the materials compared in this study.

We noted higher rates (17.9%) of implant failure in patients where β -TCP was used as an osteobiologic along with higher rates of pseudoarthrosis (28.2%) as noted in some preclinical studies on β -TCP which needs further investigation.³⁰

Limitations

The present systematic review has some limitations needing recognition for an adequate interpretation of the results. First,

the heterogeneity in the included studies concerning the study designs and graft usage prevented us from giving strong recommendations based on the results obtained. We did not analyze the complications such as donor site complications and subsidence of the grafted material which might mask the real incidence of the complications analyzed. We did not include studies that used these grafts without any report of complications, since our primary objective is to identify the rate of occurrence of each complication following ACDF surgery.

Authors Perspective

Upon analyzing the studies for the osteobiologics to aid in fusion in ACDF surgery, compared to the autografts, none of the included materials such as DBM or β -TCP, or HA had an evident rise in immediate complications such as dysphagia, dysphonia, neck pain, neurological worsening, infection or dural tear. Hence, their immediate safety is well established. However, complications such as implant failure, and pseudoarthrosis were predominantly noted in fusion constructs using β -TCP while ASD is predominantly noted in fusion constructs using DBM. Hence, future studies are warranted to analyze the material properties of β -TCP and DBM on their suitability for use in ACDF fusion constructs. We stress that the results were all derived from studies with limited methodological robustness and caution should be exercised before considering the results for clinical translation.

Conclusions

Degenerative cervical conditions treated with one or two-level ACDF surgery using DBM, HA, or β -TCP with or without cervical plating are associated with complications such as adjacent segment disease, dysphagia, and pseudoarthrosis. However, consequent to the study designs and clinical heterogeneity of the studies, it is not possible to correlate these complications accurately with any specific graft material employed. Further well-designed prospective studies are needed to correctly know the related morbidity of each graft used for achieving fusion in ACDF.

Declaration of Conflicting Interests




The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: ATEC-attended course in San Diego, travel and lodging supported by ATEC, ATEC-own stock options that were purchased personally. Fehling Instruments GmbH, Stayble Therapeutics Via Regenerate Life Sciences GmbH, Mundipharma Via Regenerate Life Sciences GmbH, Spinplant GmbH. Depuy Synthes. Yes, various lawfirms me, AO Foundation Board reimbursements, Bone biologics Personal investment, Surgitech Personal investment, Pearldiver Personal investment. SeaSpine (past, paid to the institution), Next Science (paid directly to institution), Motion Metrics (past, paid to the

institution), NIH SBIR grant NIH SBIR Subaward, MAX Bio-Pharma, Cerapedics (past), The Scripps Research Institute (past), AO Spine (past), Xenco Medical (past), AO Spine, North American Spine Society, US 2014O194364A1, US010391142B2, North American Spine Society: committee member, Lumbar Spine Society: Co-chair Educational Committee, AOSpine Knowledge Forum Degenerative: Associate member, AOSNA Research committee-committee member. Medtronic, Baxter, SafeOrthopaedics, Croatian Society of Vertebratology, Medtronic, Paid consultant for company, SI Bone Research support to institution, AO Spine Research support to institution, Mizuho Orthopedics Systems Research support to institution, Medtronic Paid consultant for company.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This supplement was organized and financially supported by AO Spine through the AO Spine Knowledge Forum Degenerative, a focused group of international spine experts.

ORCID iDs

Juan P. Cabrera  <https://orcid.org/0000-0003-4685-6106>
 Sathish Muthu  <https://orcid.org/0000-0002-7143-4354>
 Mohamed Kamal Mesregah  <https://orcid.org/0000-0002-8047-9159>
 Ricardo Rodrigues-Pinto  <https://orcid.org/0000-0002-6903-348X>
 Yabin Wu  <https://orcid.org/0000-0002-1836-7674>
 Hans Jörg Meisel  <https://orcid.org/0000-0003-3838-1489>
 Zorica Buser  <https://orcid.org/0000-0002-5680-0643>

Supplemental Material

Supplemental material for this article is available online.

References

- Martin C, D'Oro A, Buser Z, et al. Trends and costs of anterior cervical discectomy and fusion: A comparison of inpatient and outpatient procedures. *Iowa Orthop J*. 2018;38:167-176.
- Lebude B, Yadla S, Albert T, et al. Defining "complications" in spine surgery neurosurgery and orthopedic spine surgeons' survey. *J Spinal Disord Tech*. 2010;23(8):493-500.
- Pollock R, Alcelik I, Bhatia C, et al. Donor site morbidity following iliac crest bone harvesting for cervical fusion: A comparison between minimally invasive and open techniques. *Eur Spine J*. 2008;17(6):845-852. doi:10.1007/s00586-008-0648-3
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097
- Kirshblum S, Snider B, Rupp R, Read MS. Updates of the international standards for neurologic classification of spinal cord injury: 2015 and 2019. *Phys Med Rehabil Clin N Am*. 2020; 31(3):319-330. doi:10.1016/j.pmr.2020.03.005
- Sterne JAC, Savović J, Page MJ, et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ*. 2019;366:14898. doi:10.1136/bmj.14898
- Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological index for non-randomized studies (MINORS): Development and validation of a new instrument. *ANZ J Surg*. 2003;73(9):712-716.
- Hacker RJ, Cauthen JC, Gilbert TJ, Griffith SL. A prospective randomized multicenter clinical evaluation of an anterior cervical fusion cage. *Spine*. 2000;25(20):2646-2654.
- Agrillo U, Mastronardi L, Puzilli F. Anterior cervical fusion with carbon fiber cage containing coralline hydroxyapatite: Preliminary observations in 45 consecutive cases of soft-disc herniation. *J Neurosurg*. 2002;96(3 suppl 1):273-276.
- Cauthen JC, Theis RP, Allen AT. Anterior cervical fusion: A comparison of cage, dowel and dowel-plate constructs. *Spine J*. 2003;3(2):106-117.
- Cosar M, Ozer AF, Celal Iplikcioglu A, et al. The results of b-tricalcium phosphate coated hydroxyapatite (b-TCP/HA) grafts for interbody fusion after anterior cervical discectomy. *J Spinal Disord Tech*. 2008;21(6):436-441.
- Akula M, Taha M, Mathew B, O'Reilly G. The plate cage benezech implant as an alternative to autologous bone graft in the treatment of cervical spondylosis: Clinical and functional outcome. *Br J Neurosurg*. 2008;22(4):542-545. doi:10.1080/02688690802057326
- Moon HJ, Kim JH, Kim JH, Kwon TH, Chung HS, Park YK. The effects of anterior cervical discectomy and fusion with stand-alone cages at two contiguous levels on cervical alignment and outcomes. *Acta Neurochir*. 2011;153(3):559-565. doi:10.1007/s00701-010-0879-z
- Niu C-C, Liao J-C, Chen W-J, Chen L-H. Outcomes of interbody fusion cages used in 1 and 2-levels anterior cervical discectomy and fusion titanium cages versus polyetheretherketone (PEEK) cages. *J Spinal Disord Tech*. 2010; 23(5):310-316.
- Zagra A, Zagra L, Scaramuzza L, Minoia L, Archetti M, Giudici F. Anterior cervical fusion for radicular-disc conflict performed by three different procedures: Clinical and radiographic analysis at long-term follow-up. *Eur Spine J*. 2013;22 Suppl 6(suppl 6): S905-S909. doi:10.1007/s00586-013-3006-z
- Kim CH, Chung CK, Hahn S. Autologous iliac bone graft with anterior plating is advantageous over the stand-alone cage for segmental lordosis in single-level cervical disc disease. *Neurosurgery*. 2013;72(2):257-265. doi:10.1227/NEU.0b013e31827b94d4
- Shin JS, Oh SH, Cho PG. Surgical outcome of a zero-profile device comparing with stand-alone cage and anterior cervical plate with iliac bone graft in the anterior cervical discectomy and fusion. *Korean J Spine*. 2014;11(3):169-177. doi:10.14245/kjs.2014.11.3.169
- Kim S, Kim S. Anterior cervical discectomy and fusion using a double cylindrical cage versus an anterior cervical plating system with iliac crest autografts for the treatment of cervical degenerative disc disease. *J Korean Neurosurg Soc*. 2014;55(1):12-17.

19. Xie Y, Li H, Yuan J, Fu L, Yang J, Zhang P. A prospective randomized comparison of PEEK cage containing calcium sulphate or demineralized bone matrix with autograft in anterior cervical interbody fusion. *Int Orthop*. 2015;39(6):1129-1136. doi:10.1007/s00264-014-2610-9
20. Yi J, Lee GW, Nam WD, et al. A prospective randomized clinical trial comparing bone union rate following anterior cervical discectomy and fusion using a polyetheretherketone cage: Hydroxyapatite/B-Tricalcium phosphate mixture versus hydroxyapatite/demineralized bone matrix mixture. *Asian Spine J*. 2015;9(1):30-38. doi:10.4184/asj.2015.9.1.30
21. Chin KR, Pencle FJR, Seale JA, Valdivia JM. Soft tissue swelling incidence using demineralized bone matrix in the outpatient setting. *World J Orthop*. 2017;8(10):770-776. doi:10.5312/wjo.v8.i10.770
22. Farrokhi MR, Nikoo Z, Gholami M, Hosseini K. Comparison between acrylic cage and polyetheretherketone (PEEK) cage in single-level anterior cervical discectomy and fusion. *Clin Spine Surg*. 2017;30(1):38-46. doi:10.1097/BSD.0000000000000251
23. Yu J, Ha Y, Shin JJ, et al. Influence of plate fixation on cervical height and alignment after one- or two-level anterior cervical discectomy and fusion. *Br J Neurosurg*. 2018;32(2):188-195. doi:10.1080/02688697.2017.1394980
24. Shiban E, Nies M, Kogler J, et al. No correlation between radiological and clinical outcome 1 year following cervical arthrodesis. *Acta Neurochir*. 2018;160(4):845-853. doi:10.1007/s00701-018-3495-y
25. Cheung ZB, Gidumal S, White S, et al. Comparison of anterior cervical discectomy and fusion with a stand-alone interbody cage versus a conventional cage-plate technique: A systematic review and meta-analysis. *Global Spine J*. 2019;9(4):446-455. doi:10.1177/2192568218774576
26. Oshina M, Oshima Y, Tanaka S, Riew KD. Radiological fusion criteria of postoperative anterior cervical discectomy and fusion: A systematic review. *Global Spine J*. 2018;8(7):739-750. doi:10.1177/2192568218755141
27. Rhee JM, Chapman JR, Norvell DC, Smith J, Sherry NA, Riew KD. Radiological determination of postoperative cervical fusion: A systematic review. *Spine*. 2015;40(13):974-991. doi:10.1097/BRS.0000000000000940
28. Iunes EA, Barletta EA, Belsuzarri TAB, et al. Pseudarthrosis in anterior cervical discectomy and fusion with a self-locking, stand-alone cage filled with hydroxyapatite: A retrospective study with clinical and radiological outcomes of 98 levels with a minimum 2-year follow-up. *J Neurosurg Spine*. 2020;33(6):717-726. doi:10.3171/2020.4.SPINE20357
29. Tetreault L, Lange SF, Chotai S, et al. A systematic review of definitions for dysphagia and dysphonia in patients treated surgically for degenerative cervical myelopathy. *Global Spine J*. 2022;12:1535-1545. doi:10.1177/21925682211035714
30. Lu H, Zhou Y, Ma Y, et al. Current application of beta-tricalcium phosphate in bone repair and its mechanism to regulate osteogenesis. *Front Mater*. 2021;8:698915. doi:10.3389/fmats.2021.698915